DIRECTIONS FOR USE:

Mix proper dosage with any amount of palatable feed that will be consumed in one serving. If possible, confine treated pets for a day or two so that the droppings can be collected and destroyed. Heavily infested animals may require a second treatment after two weeks. Consult your veterinarian for assistance in the diagnosis. treatment and control of parasites.

DOSAGE TABLE Measure Accurately

)ogs

easpoon easpoon teaspoon easpoon teaspoons aspoons

aspoons

Body Weight	Cats	j
2-1/2 lbs. or less	1/4 teaspoon	1/4
5 lbs.	1/2 teaspoon	1/2
7-1/2 lbs.	3/4 teaspoon	3/4
10 lbs.	1 teaspoon	1 te
15 lbs.		1 1/2
30 lbs.		3 te
60 lbs.		6 te

ACTIVE INGREDIENT: Piperazine (Dipiperazine Sulfate)-equivalent to 4.25 grams Piperazine Base per 100cc.



For more information on safely worming your pet, call 1-800-921-PETS (1-800-921-7387)



Consult veterinarian before using in severely debilitated animals. Do not worm animals more than twice yearly except as advised by a veterinarian. This product is effective for those roundworm stages found in the intestine and does not remove migrating larval stages. Animals should be checked periodically by a veterinarian for the presence of other intestinal parasites. Avoid freezing.

> ³⁹⁰⁷⁹03081 6

Lot no. and expiration date on bottom

FARNAM Division of Famam Companies Inc P.O. Box 34820 • Phoenix, AZ 85067-4820

©1999 Farnani Companies, Inc. Printed in U.S.A. Product No. 3081