PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Keep children and pets away from treated areas until spray has dried. Avoid contamination of food. Do not contaminate water sup-

FIRST AID

. Take off contaminated clothing. Rinse skin with plenty of water for 15-20 IF ON SKIN:

Call a poison control center or doctor for treatment advice.
Hold eye open and rinse slowly and gently with water for 15–20 minutes.
Remove contact lenses, if present; after the first 5 minutes, then continue IF IN EYES:

. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters

CONSUMER INFORMATION

This product will treat up to:

1,600 sq. ft. (approx. size of ¼ tennis court) - for 8 Ft. Oz. container size



SUPER CONCENTRATE

- 41% Concentrate Ultra Formula
- . Kills Weeds & Grasses Down To The Root
- . Kills Unwanted Vegetation
- Covers Up To 1.600 Square Feet

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS 8 FL Oz.

This product can be used to refill Hi-Yield® KILLZALL™ Ready-To-Use Weed & Grass Killer, following mixing directions on the label. Use the mixed solution within 2 weeks.

Where to use

General weed control: Use near fences, in paths, patios, sidewalks, curbs and driveways, around ornamental trees, shrubs, flower beds and plantings, around buildings. Controls weeds in brick and gravel walks. Use for trimming and edging landscapes. NOT recommended for spot weed treatment in lawns, since Hi-Yield® KILLZALL™ kills all green plants, including lawn grasses. If replacing a lawn or parts of a lawn, see For lawn renovation/replacement recommendations on this label.

Spot treatments: Direct spray on undesirable weeds and grasses in and around flower beds, ornamental trees, fruit and nut trees, grapevines, shrubs, fences, driveways and walkways. Spray when air is calm. Prevent Hi-Yield* KILLZALL™ spray from contacting the leaves, green stems, or exposed roots of plants you like. Use a cardboard or plastic shield to protect desirable plants from spray or drift. Hi-Yield* KILLZALL™ will not move in or on the soil to untreated plants. If Hi-Yield* KILLZALL™ is used to control weeds around fruit or nut trees or grapevines, wait 21 days before eating the fruit or nuts.

Landscapes: For use in preparing areas for planting of ornamentals, trees, shrubs, desert landscapes, rock gardens, flower beds or similar plantings. Treated areas can be planted 1 day after treatment

Wildlife areas site preparation: Use Hi-Yield® KILLZALL™ to prepare areas for planting wildlife food plots. Exotic or undesirable plants can be controlled with Hi-Yield® KILLZALL™ to allow for planting or recovery of native plant species. Spot treatments may be directed to selectively remove unwanted plants. Avoid spray drift to desirable plants.

For lawn renovation/replacement:

Use Hi-Yield® KILLZALL™ to kill existing vegetation (weeds and grasses and the old lawn).

Apply in Spring or Fall, when daytime temperatures are at least 60°F for best results. If the lawn is

dry, water lawn every other day for 7 days before applying Hi-Yield® KILLZALL™.

Do not mow for 7 days before or after treatment.

Use 2½ fl. ozs. per gallon of water for each 300 sq. ft. of area to treat.

Wait 7 days after application to prepare the soil by raking, leveling or rototilling. Rake up and remove loosened thatch or debris.

ake up and remove looselied match of debis

4

Work in starter fertilizer and soil amendments, if needed. Level the soil.

Sow new seed according to directions on seed package, or reinstall sod. Grass seed must have good contact with soil to germinate and grow. Use temporary mulch such as straw to conserve moisture. Keep are moist to establish new lawn. Water as needed.

To Control Brush or Vines: Apply to green and actively growing brush or vines. For plants taller than 5 feet, cut them back and spray the regrowth. Fall applications to brush may delay control to the following season. A second application may be required on hard-to-control species such as Blackberry, Kudzu, or Poison Oak.

To kill Vines growing up a pole, fence or mature tree trunk, cut vine to a height of 3 to 4 feet and spray thoroughly covering remaining vine. If vine is climbing a shrub or immature (green) tree trunk, to avoid damage to these plants, cut vine at base and treat as directed for stumps or spray regrowth. If spraying regrowth, shield shrub and green trunk from drift with a sheet of plastic or cardboard.

Stump Treatment: Stumps can be treated to prevent regrowth. Apply Hi-Yield* KILLZALLT** When

- Cut stump close to the soil surface.
- Make 4–5 nail holes into the stump.
- Flood the stump with undiluted product.

This treatment will suppress or control many types of woody brush and trees, such as Alder, Bluegum Eucalputus, Madrone, Oak, Giant Reed, Saltcedar, Sweetgum, and Tan Oak.

vegetation is actively growing or immediately (within 5 minutes) after being cut down.

Reseeding and Replanting in Treated Areas Ornamentals and Flowers: Re-seed or replant 1 day following application.

Vegetables: Wait **3 days** after spraying, before seeding or transplanting the plants listed below:

togotables. Wait o days after spraying, belove seeding or transplanting the plants listed below.					
Beans	Crenshaw Melon	Mangos Melon	Radish		
Beet Greens	Cucumber	Melons	Rhubarb		
Broccoli	Eggplant	Muskmelon	Rutabaga		
Brussels Sprouts	Endive	Mustard Greens	Spinach		
Cabbage	Garlic	Okra	Summer Squash		
Cantaloupe	Honeydew Melon	Onions	Sweet Potato		
Carrot	Honey Ball Melon	Rape Greens	Tomatillo		
Cauliflower	Horseradish	Parsley	Tomato*		

Reseeding and Replanting in Treated Areas (continued)

Ornamentals and Flowers: Re-seed or replant 1 day following application.

Vegetables: Wait 3 days after spraying, before seeding or transplanting the plants listed below:

Casaba Melon	Jerusalem Artichoke	Parsnips	Turnip
Celery	Kale	Peas	Watercress
Chard (Swiss)	Kohirabi	Pepper	Watermelon
Chicory	Leeks	Persian Melon	Winter Squash
Collards	Lentils	Potato	Yams
Corn	Lettuce	Pumpkin	
		· '	

*Note: Wait 3 days before planting Tomatoes from seed and 30 days for rooted Tomato transplants. For Tomato transplants or vegetables not listed, wait 30 days after spraying before planting into treated soil.

Small Fruits and Berries: Blackberry, Black Raspberry, Blueberry, Boysenberry, Currant, Dewberry, Alderberry, Gooseberry, Huckleberry, Loganberry, Olallieberry, Red Raspberry and Strawberry may be planting by seed, rooted cutting or transplant 7 days after spraying Hi-Yield® KILLZALL™. Wait 30 days after spraying and before planting other small fruits and berries.

Fruit and Nut Trees Caneberries and Grapevines: Allow 21 days before eating the fruits or nuts when using Hi-Yield® KILLZALL™ to control weeds around fruit or nut trees, caneberries and grapevines.

STORAGE AND DISPOSAL

Storage: Use and store this product only in its original container. Store product in a secure storage area away from sources of heat or open flame. Protect product from freezing.

Disposal: Do not reuse container. To dispose of unused product and container, securely wrap original container in several layers of newspaper and discard in trash.

Spill or Leak: In case of spill or leak, soak up with paper towels and discard in trash.

The label instructions for the use of this product reflect the opinion of experts based on field uses and tests. The directions are believed to be reliable and should be followed carefully. However, it is

6

impossible to eliminate all risks inherently associated with use of this product. Desirable plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Voluntary Purchasing Groups, Inc. All such risks shall be assumed by the user.

Voluntary Purchasing Groups, Inc. warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the Directions for Use set forth in the label, subject to the risks referred to above.

Voluntary Purchasing Groups, Inc. makes no other express or implied warranty including any other express or implied warranty of FITNESS or MERCHANTABILITY.

- 1

Distributed by Voluntary Purchasing Groups, Inc. Bonham, Texas 75418

EPA Reg. No. 67760-49-7401 EPA EST. NO. 7401-TX-1



DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Read the entire panel before using this product. Use only according to label instructions observing all precautions.

NOTE: See the CONSUMER INFORMATION section of this label for important information on protecting plants you like.

Mixing Directions: For best results use 2% fl. ozs. (5 Tbs.) per gallon of water. One gallon of water will treat approximately 300 sq. ft.

Hi-Yield® KILLZALL™ eliminates nearly all common weeds, grasses and unwanted vegetation. Add water and use along fences, in driveways, sidewalks, brick and graved walkways, next to trees and shrubs, and around buildings to control weeds and unwanted vegetation. Use Hi-Yield® KILLZALL™ for edignig land-scapes. Hi-Yield® KILLZALL™ is suited for lawn renovation or to prepare large sites for planting of flower beds, vegetable gardens or wildlife food plots. Hi-Yield® KILLZALL™ can also be used for large jobs such as lawn renovation and vegetable and flower garden preparation. For edging around lawns, lawn replacement/renovation, and for garden plot preparation. Do not use on or around edible feed or food crops.

Important: This product is a non-selective herbicide that will kill all vegetation contacted. This herbicide kills unwanted plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves down to the roots. Avoid direct spray or drift onto desirable plants, lawn grasses, shrubs and trees. Protect desirable plants and vegetation with a piece of plastic or cardboard. If desirable plants are accidently sprayed, wash plants with water immediately. Any Hi-Yield® KILLZALL™ not absorbed by the plant is broken down into natural materials. This product has no soil activity and thus will not affect nearby untreated plants.

Do not mow, cut or otherwise disturb treated vegetation for at least 7 days following application.

WEEDS AND GRASSES CONTROLLED

Annual Weeds and Grasses:

Annual Ryegrass, Annual Bluegrass, Barnyardgrass, Beggarweed, Black Medic, Blue Medic, Blue Toadflax, Brassbuttons, Blue Mustard, Bromegrass, Bur Clover, Buttercup, Cheeseweed, Chickweed, Clover, Common Groundsel, Common Lambsquarters, Common Plantain, Common Ragweed, Crabgrass, Creeping Beggarweed, Diffuse Lovegrass, Dog Fennel, Evening Primrose, False Dandelion, Fall Panicum, Fennel, Fiddleneck, Field Pennycress, Field Sandbur, Filaree, Florida Pusley, Foxtail, Garden Spurge, Goosegrass, Green Foxtail, Hemp Sesbania, Henbit, Horseweed/Marestail, Ironweed, Knotweed, Lambsquarters, Little Bittle Cress, London Rocket, Maiden Gane, Mallow, Mayweed, Mouseear Chickweed*, Oxalis*, Pennsylvania Smartweed, Pennywort, Pigweed, Prickty Lettuce, Prostrate Spurge, Puncture Vine, Purslane, Redroot Pigweed, Sandspur, Shepherdspurse, Smartweed, Smooth Cat's Ear, Smooth Pigweed, Sowthistle, Spotted Spurge, Sprangletop, Spurges, Tansy Mustard, Tansy Ragwort, Fleaweed, Texas Panicum, Yellow Oxalis, and many other annual grasses and weeds. Controls many unwanted plants and kills most types of weeds and grasses.

Perennial Weeds, Grasses and Brush (Tough weed control):

Alder, Artichoke Thistle, Bahiagrass, Barnyardgrass, Bentgrass, Bermudagrass, Blackberries, Bluegrass, Broadlear Plantain, Brownseed Paspalum, Buckhorn Plantain, Bull Thistle, Canada Thistle, Cattail, Oeanothus, Centipedgrass, Cherry, Cooklebur, Cogongrass, Common Mullein, Common Ragweed, Coralbead, Coyote Brush, Creeping Bentgrass, Creeping Charlie, Creeping Lantana, Curly Dock, Dallisgrass, Dandelion, Dewberry, Dog Fennel, Elderberry, False Dandelion, Fescue Species, Field Bindweed, Guineagrass, Hairy Crabgrass, Honeysuckle, Horsenettle, Horseradish, Iceplant, Johnsongrass, Kentucky Bluegrass, Kikuyugrass, Knapweed, Knawel, Kudzu, Lantana, Malden

Grass, Milkweed, Mouseear Chickweed*, Multiflora Rose, Kimblewill, Nutgrass (Nutsedge), Oak Species, Oldenlandia, Orchardgrass, Oxalis*, Pampasgrass, Pennywort, Perennial Ryegrass, Plantain, Poison Oak, Poison lvy, Poison Hemlock, Primrose, Purple Budweed*, Purple Nutsedge, Quackgrass, Quaking Aspen, Fragweed, Raspberry, Red Clover, Smooth Bromegrass, Sourdock, Sowthistle, St. Augustinegrass, Sumac, Tall Fescue, Tansy Ragwort, Thistle, Timothy, Torpedograss, Trumpetcreeper, Vaseygrass, Virginia Creeper, White Clover, Whitetop, Wild Barley, Wild Blackberry, Wild Carrot, Wild Morning-glory, Wild Oats, Wild Sweet Potato, Willow, Yarrow, Yellow Nutgrass, Yellow Starthistle, Zoysia and many other perennial grasses, weeds, brush and sedges. Controls many unwanted weeds/plants and kills most types of weeds and grasses.

*Annual or Perennial

When to apply:

Apply to actively growing weeds, preferably before the seed heads form. Completely spray the foliage of the weeds and grasses you want to kill until thoroughly wet. It's best to apply on a warm sunny day when daytime temperatures are above 60°F. Weed control is accelerated by warm sunny weather. Apply when the wind is calm. Reapply if it rains within 24 hours of application. Hard-to-control weeds may require a second application if not completely dead in 4 weeks.

tow to apply:

This product is formulated for "spot" treatment of individual weeds and grasses or for treatment of large areas as instructed on this label (see specific site areas below). Apply with a plastic, aluminum or stainless steel tank-type sprayer.

Do not apply with galvanized or unlined steel sprayer, or through any irrigation system or sprinkling can.

Use a tank-type sprayer with a coarse spray nozzle for spot treatments. Use hose-end sprayer to treat large areas such as for lawn renovation or to prepare sites for planting flower beds or vegetable gardens.

Apply to evenly and thoroughly wet weeds and grasses.

Apply Hi-Yield® KILLZALL™ using any of the following equipment:

- Plastic, aluminum or stainless steel tank sprayer
 A hose-end spayer recommended for weed killers
- A nose-end spayer recommended for weed k
- Hand-trigger pump sprayer

Cleaning Application Equipment:

To clean sprayer after use, triple rinse sprayer and all sprayer parts with water. Spray rinse water on

bare soil or gravel. After thorough cleaning, sprayer can be used for other products.

Directions for use:

Use the following table to determine the amount of Hi-Yield® KILZALL™ to mix with water. Close cap tightly after use. No need to rinse cap.

1 tablespoon (Tbs.) = 3 tsp.

1 teaspoon (tsp.)

2 Fl. Oz. (4 Tbs)

1 FI. UZ. (Z 1115)) 2 FI. UZ. (4 TDS)				
	MIXING IN:				
APPLICATOR	General weed control annuals and seedling weeds	Tough weed control perennials or brush or lawn renovation.	AREA TO TREAT		
Tank Sprayer	Use1½ oz. (3 Tbs) in 1 gallon of water.	Use 2½ oz. (5 Tbs.) in 1 gallon of water.	300 sq. ft. (30 ft.x 10 ft.) or use for spot treatment.		
Hose-End Sprayer	Add 1½ oz. (3 Tbs.) to sprayer jar. sprayer jar. DO NOT ADD WATER. Set dial to 1½ oz. and spray.	Add 2½ ozs. (5 Tbs) to (30 ft. x 10 ft.) DO NOT ADD WATER. Set dial to 2½ oz. and spray.	300 sq. ft.		
6 Gallon Hose- End Sprayer	Add 6 oz. to sprayer jar. Fill to the 4 gallon mark and spray.	Add 10 oz. to sprayer jar. Fill to the 4 gallon mark and spray.	1200 sq. ft. (30 ft.x40 ft.)		
Hand trigger sprayers 24 oz. capacity	Use 2 tsp in 24 oz. of water.	Use 1 Tbs in 24 oz. of water.	Use for spot treatments or apply to single		

2