

FIRST AID

If Swallowed	<ul style="list-style-type: none"> • Call Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
If In Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a Poison Control Center or doctor for treatment advice.
If On Skin Or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact the National Pesticide Telecommunications Network at 1-800-858-7378 for emergency medical treatment.

Note to Physician

Probable mucosal damage may contraindicate gastric lavage.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER: Fatal if swallowed, inhaled or absorbed through skin. This product is corrosive to flesh because of its caustic alkaline nature, and it causes irreversible eye damage and skin burns. Do not breathe dust, vapor, spray mist, or get in eyes, or get on skin or clothing. Wear long-sleeved shirt and long pants, socks, chemical-resistant footwear, chemical resistant gloves made of any waterproof material, safety glasses or goggles and hat. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur.

PHYSICAL AND CHEMICAL HAZARDS

May give off highly toxic and extremely flammable hydrogen sulfide gas if mixed with acids.

Hi-Yield®

Improved

Lime Sulfur Spray



- Fungicide, Insecticide, Miticide For Fruits, Nuts, Ornamentals, Roses.

**KEEP OUT OF REACH
OF CHILDREN**

DANGER PELIGRO

Si usted no entiende la etiqueta,
busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label,
find someone to explain it to you in detail.)

SEE ADDITIONAL PRECAUTIONARY STATEMENTS INSIDE
BOOKLET. ALL APPLICABLE DIRECTIONS, RESTRICTIONS
AND PRECAUTIONS ON THE LABEL ARE TO BE FOLLOWED.
SEE DIRECTIONS FOR USE IN BOOKLET.

NET CONTENTS ONE QUART (32 FL. OZS.)

ACTIVE INGREDIENTS:

Calcium Polysulfides 28%

INERT INGREDIENTS:..... 72%

TOTAL:100%

PLUMS: *Blotch, Powdery Mildew.* Apply 4 teaspoons per gallon of water. Begin 3 to 5 weeks before Harvest. Repeat weekly until picking. *Black Knot.* Apply 8 fl. oz. per gallon of water. Apply in green tip stage. *Brown Rot (Fruit Rot).* Apply 1 fl. oz. per gallon of water. Apply 20, 10 and 2 days before harvest. *Leaf Curl.* Apply 12½ to 15 fl. oz. per gallon of water. Apply in dormant stage.

PRUNES: *Black Knot.* Apply 8 fl. oz. per gallon of water. Apply in green tip stage. *Brown Rot (Fruit Rot), Leaf Spot, Powdery Mildew.* Apply 4 teaspoons per gallon of water and begin 3 to 5 weeks before Harvest. Apply at weekly intervals until picking.

RASPBERRIES: *Anthraxnose, Powdery Mildew.* Apply 10½ fl. oz. per gallon of water. Spray once when new canes are 8 to 12 inches long. *Cane Blight.* Apply 1¼ fl. oz. per gallon of water. Apply at Spring bud swell. Repeat in late Fall dormant period. *Spur Blight.* Apply 12½ to 15 fl. oz. per gallon of water. Apply in delayed dormant stage when buds begin to break and show silver.

SHADE TREES, SHRUBS: For control of *San Jose Scale, Leaf Blotch, Olive Scale,* and as an aid in the control of *Powdery Mildew, Anthracnose.* Apply 5 fl. oz. plus 1 fl. oz. light medium Summer Oil per gallon of water once in early Spring before buds swell. Do not apply to Evergreens, *Euonymus* or *Rhododendrons* or allow spray to drift on these susceptible species. Full Season control of *Powdery Mildew* and *Anthraxnose* will require additional application of Fungicide. Keep agitated during spraying.

EARLY SPRING SPRAYING (AS GROWTH STARTS)

ROSES: *Rose and San Jose Scale, Powdery Mildew.* Apply 4 fl. oz. per gallon of water. Apply in Spring before buds swell and again in Fall after leaves drop.

SUMMER FOLIAGE SPRAYING

ROSES: *Black Spot, Powdery Mildew, Rust, Red Spider Mites.* Apply ½ fl. oz. per gallon of water. Apply at 10 to 15 day intervals in Growing Season.

COMBINATION WINTER SPRAY

FRUIT TREES, ROSES AND DECIDUOUS HEDGE PLANTS, SHADE TREES AND SHRUBS (DORMANT AND DECIDUOUS ONLY): *San Jose Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, Overwintering Insect Eggs, Many Overwintering Fungus Spores, Or Plant Disease.* Apply 4 fl. oz. plus 1¼ fl. oz. dormant spray oil per gallon of water. Apply one dose as Winter buds swell before opening. Keep agitated during spraying.

TABLE OF EQUIVALENTS FOR HOME GARDEN ONLY

VOLUME

1 Fluid oz. = 2 Tablespoons = 6 Teaspoons • 1 Gallon = 4 Quarts = 8 Pints

CONDITIONS OF SALE LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of Voluntary Purchasing Groups, Inc. or Seller. In addition, failure to follow the label directions may cause injury to crops, animals, man or the environment. Voluntary Purchasing Groups, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions For Use subject to the factors noted above which are beyond the control of Voluntary Purchasing Groups, Inc. Voluntary Purchasing Groups, Inc. makes no other warranties or representations of any kind, express or implied concerning the product, including no implied warranty of merchantability or fitness for any particular purpose. The exclusive remedy against Voluntary Purchasing Groups, Inc. for any cause of action relating to the handling or use of this product is a claim of damage and in no event shall damages or any other recovery of any kind against Voluntary Purchasing Groups, Inc. exceed the price of the product, which causes the alleged loss, damage, injury or other claim. Voluntary Purchasing Groups, Inc. shall not be liable and any and all claims against Voluntary Purchasing Groups, Inc. are waived, for special, indirect, incidental, or consequential damages or expense, of any nature, including, but not limited to, loss of profits, or income, whether or not based on Voluntary Purchasing Groups, Inc. negligence, breach of warranty, strict liability in tort or any other cause of action. Voluntary Purchasing Groups, Inc. and the Seller offer this product and the Buyer and User accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

VOLUNTARY PURCHASING GROUPS, INC.
P. O. BOX 460
BONHAM, TEXAS 75418
EPA Reg. No. 71096-6-7401
EPA Est. No. 7401-TX-1



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. For use on Plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or golf courses or lawns and grounds. Do not enter treated areas until sprays have dried.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE LABEL ARE TO BE FOLLOWED.

Faulty spray equipment, highly concentrated materials, or extremes of weather during or following spraying may lead to fruit or foliage injury. The risk of spray injury is greater when drought stress exists. The user is advised not to use Lime Sulfur on any crop unless local use has proved that Lime Sulfur does not damage crops in that locality.

Do not use Lime Sulfur on Apricots. Do not use Oil with Lime Sulfur in Summer applications. Do not apply when temperature exceeds 85°F. Do not apply Oil following Lime Sulfur, nor Lime Sulfur following Oil, in foliage period. Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

Storage: Store product in a secure locked place, inaccessible to children, pets, and livestock. Store in a cool, dry place. Keep container closed when not in use.

Container Disposal: If Empty: Do not reuse this container. Place in trash or offer for recycling, if available. **If Partly Filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE DIRECTIONS

Spray to cover foliage until wet. Spray with the wind to your back. If spray contacts concrete or white, painted surfaces, discoloration of this surface may occur.

ALMONDS: Powdery Mildew, Shothole. Apply 10½ fl. oz. per gallon of water. Apply every 2 weeks between November 15 and December 15. Dormant application only.

APPLES: Blotch. Apply 2½ fl. oz. per gallon of water. Apply in pre-bloom and petal fall stages and 10 days later. **Leaf Spot.** Apply 9½ to 13 fl. oz. per gallon of water in delayed dormant stage or 4 teaspoons per gallon of water in growing season. **Powdery Mildew.** Apply 3.2 fl. oz. per gallon of water. Apply in pre-pink, pink and calyx stages. **Scab.** Apply 2 to 2½ fl. oz. per gallon of water in delayed dormant, pre-pink and pink stage. Caution: Use on Delicious Apples may result in injury. No time limitation.

BLACKBERRIES: Anthracnose, Powdery Mildew, Rust. Apply 7½ to 15 fl. oz. per gallon of water. Apply in dormant (green bud) stage. Use 4 teaspoons per gallon water when fruiting canes are 8 to 12 inches long and before blossoms have opened. **Overwintering Fungus Spores, Cane Blight.** Apply 3 fl. oz. per gallon of water. Apply in dormant stage.

BLUEBERRIES: (Registration and use limited to Pacific Northwest) To aid control of **Mummy Berry**, (caused by the fungus Sclerotinia Urnula), apply a post-harvest (September or October) or early dormant spray of 10¼ fl. oz. per gallon of water.

CHERRIES (SWEET): Powdery Mildew. Apply 9½ fl. oz. per gallon of water. Apply in delayed dormant stage.

CHERRIES (TART): Brown Rot Blossom Blight, Leaf Spot, Scab. Apply 2 fl. oz. per gallon of water. Apply in pre-bloom, early bloom and full bloom stages. **Brown Rot** (Fruit). **Leaf Spot, Powdery Mildew.** Apply 4 teaspoons per gallon of water. Apply 3 to 5 sprays at weekly intervals just before Harvest.

CITRUS: Red Spider Mites. Apply 2½ fl. oz. per gallon of water. Apply when the Mites appear. **Thrips.** Apply 2½ fl. oz. per gallon of water. Apply as recommended by your State Agricultural Experiment Stations.

CURRENTS, GOOSEBERRIES: Anthracnose. Apply 3¼ fl. oz. per gallon of water. Apply at bud break stage, and 10 to 15 days later. Use ½ fl. oz. per gallon of water at 10 day intervals after second spray if necessary. **Powdery Mildew.** Apply ½ fl. oz. per gallon of water. Apply before and after bloom.

FRUIT TREES (BEARING): Overwintering Fungus Spores. Apply 4½ fl. oz. per gallon of water. Apply in dormant stage.

FRUIT TREES (NON-BEARING): Blotch, Powdery Mildew, Scab. Apply 5 teaspoons per gallon of water. Apply at cluster bud and pre-bloom stages.

GRAPES: Anthracnose, Powdery Mildew. Apply 4 teaspoons per gallon of water. Apply when new shoots are 4 to 6 inches long. **(Dormant only): Powdery Mildew, Mealybugs.** Apply 2½ to 6½ fl. oz. per gallon of water.

NECTARINES: Brown Rot, Leaf Spot, Shothole. Apply 4 teaspoons per gallon of water. Apply 3 to 5 times at weekly intervals before Harvest. **Leaf Curl.** Apply 7¾ fl. oz. per gallon of water. Apply in early Winter and late dormant period. **Powdery Mildew.** Apply 4 to 6 teaspoons per gallon of water. Apply when Disease appears.

PEACHES: Brown Rot, Leaf Spot. Apply 4 teaspoons per gallon of water. Apply 3 to 5 times at weekly intervals before Harvest. **Brown Rot Blossom Blight.** Apply 1 to 2 fl. oz. per gallon of water. Apply in pre-bloom, early bloom and full bloom stages. **Leaf Curl, Scab, Shothole.** Apply 12½ to 15 fl. oz. per gallon of water. Apply in dormant season. **Powdery Mildew.** Apply 9½ fl. oz. per gallon of water at green tip stage. Use ½ fl. oz. per gallon of water on Summer foliage.

PEARS: Pear Psylla, Rust Mite, Blister Mite. Apply 5 fl. oz. per gallon of water plus 1 fl. oz. light medium Summer Oil. Apply as a Post-Harvest spray. **Bud Mite** (Pacific Coast States). Apply 6½ fl. oz. plus 2½ fl. oz. light medium Summer Oil. Apply in the Fall after temperature cools but before the first frost at the time mites first penetrate under the bud scales. **Pear Leaf Blister Mite.** Apply 6½ fl. oz. to 9½ fl. oz. per gallon of water. Apply in dormant season up to time buds swell. May be combined with oils labeled for dormant use. **Scab.** Apply 6½ fl. oz. per gallon of water applied at cracked bud stage and 4 to 6 teaspoons to 1 gallon of water at cluster bud stage.