

Prescription Treatment[®] brand **PERMA-DUST[®]**

Pressurized Boric Acid Dust

KILLS: Ants, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Firebrats, Flour Beetles, Grain Weevils, Ground Beetles, Indian Meal Moths, Millipedes, Pillbugs, Scorpions, Silverfish, Sowbugs, Spiders, and Trogodermas.

FOR USE IN AND AROUND: Apartments, Homes, Hospitals, Hotels, Food Storage Areas, Inedible Product Areas of Meat Packing Plants, Food Processing Plants, Motels, Nursing Homes, Restaurants and Other Food Handling Establishments, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Planes[†]) and Warehouses and Other Commercial and Industrial Buildings.

[†] Do not use in aircraft cabins.

ACTIVE INGREDIENTS:

Boric Acid 35.5%

INERT INGREDIENTS:

Contains petroleum distillates. 64.5%

EPA Reg. No. 499-384 EPA Est. No. 499-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

Special precautions should be taken so that all residue will be out of reach of children or domestic animals. Avoid contamination of food and foodstuffs. See side panel for additional precautionary statements.

SHAKE WELL BEFORE USING

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Before use shake until agitator ball rattles. For best results, Use in Upright Position Only. Do not spray on plastic, painted or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, telephones, etc. where the powdery residue may foul or short electric contacts and circuits.

HOW TO USE WHITMIRE MICRO-GEN CRACK AND CREVICE[®] INJECTION SYSTEM:

Use Whitmire Micro-Gen Perma-Dust[®] Pressurized Boric Acid Dust with the supplied actuator and injection tubes. Inject Perma-Dust[®] into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. Move injector tip along cracks while treating at the rate of three linear feet per second. For closed voids calculate the void's cubic area and

treat at the rate of three to five seconds per three cubic feet. For closed void areas drill small access holes. Several holes may be required in long-running voids. Repeat as necessary. This product can leave a white residue. This residue can normally be removed with a damp cloth or sponge.

INDOOR TREATMENTS

CRAWLING INSECTS - Centipedes, Cockroaches, Crickets, Dermestids, Earwigs, Millipedes, Scorpions, Sowbugs and Spiders: Inject with the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewer, floor drains and meter boxes. Repeat as necessary.

Booklice, Firebrats & Silverfish: Inject into cracks and crevices associated with baseboards, door frames, bookcases and other harborage sites. Repeat as necessary.

Ants: Inject cracks, crevices and voids where ants may be traveling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices and void areas.

STORED PRODUCT PESTS (Adult and Larval Stages) - Chocolate Moths, Drugstore Beetles, Flour Beetles, Grain Weevils, Indian Meal Moths, Trogoder-mas:

Inject into cracks and crevices and/or voids where these insects may be harboring. Infested areas include, but are not limited to, cracks and crevices in cabinets, pantries, wall voids and in machinery. Repeat treatment as necessary.

Carpenter Bees: Locate bee entrance holes. Inject 5 - 10 seconds of product into entrance holes. Seal entrances after application.

APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS

Places other than private residences in which food is held, processed, prepared or served.

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE[®] TREATMENT ONLY:

Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) also would be considered a food area. Apply directly into cracks and crevices and void areas using injection tube in points

between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD-HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE® TREATMENT ARE NOT PERMITTED.

INDOOR AND OUTDOOR TREATMENTS

Carpenter Ants: Locate source of infestation or areas of suspected activity. Inject 5-10 seconds of product into insect tunnels and cavities. Treat points of entry or exits from voids such as around doors, windows and siding.

Boxelder Bugs, Clover Mites, Cluster Flies and Elmleaf Beetles: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames. Repeat as necessary.

OUTDOOR TREATMENTS

Ants, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Ground Beetles, Millipedes, Pillbugs, Scorpions, Sowbugs

and Spiders. Treat cracks around doors, windows, under siding and voids foundation level, beneath guards on loading docks, into cracks in foundations and floors and in other places where these pests may enter building. Repeat application as necessary.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Replace cap and discard container in trash, sanitary landfill or by other approved state and local procedures. Do not incinerate or puncture.

PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, apply artificial respiration, preferably mouth-to-mouth. Get medical attention.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Apply this product only as specified on the label. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat, sparks or open flame. Contents under pressure. Do not puncture or incinerate container. Exposure to temperature above 130° F may cause bursting.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



A Prescription Treatment® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.

3568 Tree Court Industrial Blvd.

St. Louis MO 63122-6682

www.wmmg.com

© 1998 Whitmire Micro-Gen Research Laboratories, Inc.



PERMA-DUST® Pressurized Boric Acid Dust